

Articles

Adverse Effects of Topical Corticosteroid Use

DAVID A. FISHER, MD, *Antioch, California*

Topical corticosteroid use, a common and often efficacious therapy for a wide variety of cutaneous conditions, may have substantial adverse effects. These range from the notable nondermatologic side effects of hypothalamic-pituitary-adrenal axis suppression, Cushing's disease, femoral head osteonecrosis, and cataracts to a variety of less serious skin effects such as cutaneous tinea and contact dermatitis. The broad availability, efficacy, relative low cost, and ease of applying topical corticosteroids should not induce complacency or a cavalier attitude in prescribers. Physicians should have the same awareness of the possible side effects of topical steroid use as when prescribing parenteral medication.

(Fisher DA: Adverse effects of topical corticosteroid use. *West J Med* 1995; 162:123-126)

The introduction of the ultrahigh-potency topical corticosteroids has proved of important benefit for some hitherto corticoid-resistant dermatoses, but as in every new development in medicine, an adverse side to this therapeutic advance exists.

Fortunately, systemic side effects of topical corticosteroids are rare, but may occur, especially in infants¹ and elderly patients.^{2(p81)} Possible systemic side effects are related directly to such factors as the application site, duration of application, potency, and occlusion of the medication.² The application of high-potency corticosteroids should be limited, when possible, to a twice-a-day basis for three to four weeks.³ Longer and more frequent application of topical corticosteroids carries the risk of suppressing the hypothalamic-pituitary-adrenal axis and even producing Cushing's syndrome, especially in very young patients.^{4,6} Physicians must also be aware of increased transcutaneous penetration in certain areas of the body with thin epidermis, such as the eyelids, periorbital area, axillas, crural region, and genitalia.⁷ A patient with an integumentary perturbation, the quintessential example being exfoliating dermatitis, has a much greater amount of percutaneous penetration.⁸ A recent report of topical clobetasol propionate-induced femoral head osteonecrosis describes additional serious systemic effects of topical high-potency corticosteroid use.⁹

Local side effects, unlike systemic ones, are relatively frequent and becoming even more so with the introduction of the ultrahigh-potency topical corticosteroids.¹⁰ Striae distensae may develop, especially in the groin and axillas, with the prolonged application of even medium-potency corticosteroids (Figure 1).¹¹

Clotrimazole and betamethasone dipropionate (Lotrisone), which can be an effective topical agent for treating intertrigo, with long-term use has the potential of rapidly inducing striae, especially in susceptible adolescents and younger adults when used more than seven to

ten days in the crural, axillary, and flexural regions (Figure 2 and Figure 3). Another adverse effect that occurs more often in older persons and in young children, but that can occur in all age groups, is cutaneous atrophy with telangiectasia and stellate pseudoscars, especially with the prolonged use of the high- and ultrahigh-potency topical corticosteroids (Figures 4, 5, and 6).^{12,13} Adverse effects of topical corticosteroid use on the supporting collagen stroma, by inhibiting collagen production of fibroblasts, may lead to the occurrence of purpura and induce or aggravate Bateman's senile purpura.¹⁴ Clinicians generally should eschew the use of anything but low-potency corticosteroids on the face because of the rapid induction of erythema, corticoid rosacea, and perioral and periocular acneiform dermatoses (Figure 7).¹⁵

Unfortunately, the pharmaceutical industry has not always been vigilant in warning physicians concerning these possible adverse effects. An example is with mometasone furoate (Elocon), which was promoted as being safe for facial conditions, but in reality is a medium-potency steroid. Succumbing to a representative's enthusiastic sales pitch for the medication and assurances that it was safe for facial use, I produced a case of this syndrome. I have since managed several cases of mometasone-induced steroid rosacea. After complaints from me and others, the company now states that it is safe to use only for two weeks for facial dermatoses.

I strongly entreat clinicians to minimize the use of medium- and high-potency corticoids on the face of any patient. The lower-potency corticosteroids such as desonide, and even 2% to 0.5% hydrocortisone, should have adequate anti-inflammatory therapeutic benefit without the risk of producing often-difficult-to-manage steroid rosacea and erythema. Indeed, a problem with a medium- and higher-potency corticosteroid on the face is that it induces rebound erythema in which the topical corticosteroid initially causes vasoconstriction, but when



Figure 1.—Striae occur in the groin due to the use of topical clotrimazole and betamethasone dipropionate (Lotrisone).

withdrawn, produces vasodilation higher than the baseline in a trampoline-like manner.¹⁵ I have found a course of tetracycline or erythromycin, perhaps a short burst of intermittent parenteral corticosteroids, and time to be the only approach that benefits this most distressing condition. Topical corticosteroids can facilitate the proliferation of *Propionibacterium acnes*, and perhaps *Demodex folliculorum*, leading to an acne rosacea-like condition (Figure 7) that, fortunately, is often amenable to parenteral antibiotics coupled with topical keratolytic and antibacterial agents.^{16,17} The long-term application of topical corticosteroids can induce milia in older patients.¹⁸

A prolonged use of topical corticosteroids on the eyelids can not only cause thinning of the eyelids but also induce open-angle glaucoma and cataracts from transpalpebral tarsal penetration.¹⁹ Topical corticosteroids may suppress the normal cutaneous immune response to dermatophytes and thus enhance fungal infections. This can present an atypical clinical picture, to which the rubric tinea incognito has been applied.²⁰ I have observed cases of tinea in which betamethasone, a valuable corticosteroid moiety of Lotrisone, seemed to have a dominant effect over the antifungal aspect of the clotrimazole and worsened rather than cleared the cutaneous tinea. Cutaneous dermatophyte infections



Figure 2.—Striae and skin breakdown due to the use of topical medium-potency corticosteroids are shown.

TABLE 1.—Possible Adverse Effects of Topical Corticosteroid Use

Systemic Absorption	
Hypothalamic-pituitary-adrenal axis suppression	
Cushing's disease	
Femoral head osteonecrosis	
Local Effects	
Striae distensae	Milia
Cutaneous atrophy	Cataracts
Stellate pseudoscars	Tinea
Telangiectasia	? Candidiasis
Purpura	Scabies
Erythema	Hypertrichosis
Rosacea	Hypopigmentation
Acneiform dermatoses	Contact dermatitis
Rebound erythema	Tachyphylaxis
Demodicidosis	

should be treated only with a topical fungicide. A clinical study of 176 patients with potassium hydroxide and fungal culture-proven tinea pedis unequivocally documents the superiority of a single antifungal topical agent, in this instance naftifine hydrochloride, over the antifungal corticosteroid combination of clotrimazole and betamethasone propionate.²¹ The decision whether to use oral antifungal therapy is a matter of clinical judgment. Topical corticosteroid use may promote cutaneous candidiasis, although experimental proof of this is lacking.²² Local steroid use may induce atypical or extensive crusted scabies.²³ Hypertrichosis is a rare side effect of the long-term use of topical steroids. When it does occur, it is most prevalent on the face and ears.²⁴ Hypopigmentation from high- and ultrahigh-potency steroids is a possible consequence when used on a dark-skinned person.²⁵

Reports of cases of type IV allergic contact dermatitis due to local steroid use are increasing.²⁶ The rash may be chronic, erythematous, edematous, papular, and at times scaly, as opposed to the usual microvesicular eczema of acute contact dermatitis. This somewhat alarming development has not been totally elucidated,



Figure 3.—Higher-potency topical corticosteroid use produced striae on the arms.



Figure 4.—Cutaneous atrophy is shown from intralesional corticosteroid therapy.

but is the cause of some cases of corticosteroid-resistant dermatitis.²⁷ Unfortunately, the screening corticosteroids tixocortol pivalate and budesonide are unavailable for patch tests in the United States because of the Food and Drug Administration (FDA) requirement that allergens for patch testing are a medication and must be FDA-approved.²⁸ The use of these chemicals does not identify all cases of corticosteroid hypersensitivities, but according to a recent report, it identifies more than 80% of them.²⁹

Practitioners should also be mindful of the possibility of allergic contact dermatitis due to certain excipients of varying concentrations in topical corticosteroid preparations. These may be type IV delayed-hypersensitivity sensitizers, such as lanolin, paraben, and quaternium 15, or chemicals capable of producing an immediate contact urticaria, either on a nonimmunologic histamine-releasing basis or from an immunoglobulin E-mediated, type I immediate reaction.³⁰ The preservatives sorbic acid, benzoic acid, and formaldehyde are examples of such contact urticaria inducers.³¹

Topical corticosteroids may induce tachyphylaxis with extended use.³² This occurs most often in patients with psoriasis. This is another reason why it is



Figure 5.—Higher-potency topical corticosteroid use produced cutaneous atrophy with resultant skin fragility and erosions.



Figure 6.—Telangiectasia of the nose has occurred from the use of topical medium-potency corticosteroids.

recommended that the frequency of the application of ultrahigh-potency topical corticosteroids be reduced after the first two to three weeks to no more than four or five times a week.

Physicians are always mindful of the possible adverse systemic effects of the use of parenteral medications. We should be no less mindful of the possible undesirable side effects of this therapeutically valuable topical medicament.

Acknowledgment

Howard I. Maibach, MD, Professor of Dermatology, University of California, San Francisco, School of Medicine, reviewed the manuscript and made many helpful suggestions. Richard Odom, MD, provided the photographs for Figures 3, 4, and 5.

REFERENCES

1. Nachman RL, Esterly NB: Increased skin permeability in pre-term infants. *J Pediatr* 1971; 75:628-632
2. Wester RC, Maibach HI: In vivo percutaneous penetration, chap 4. *In* Marzulli FN, Maibach HI (Eds): *Dermatotoxicology*, 4th edition. Washington, DC, Hemisphere Publishing, 1991, pp 75-96
3. Olsen EA, Cornell RC: Topical clobetasol-17-propionate—Review of its clinical efficacy and safety. *J Am Acad Dermatol* 1986; 15:245-255
4. Pascher F: Systemic reactions to topically applied drugs. *Int J Dermatol* 1978; 17:768-775
5. Keipert JA, Kelly R: Temporary Cushing's syndrome from percutaneous absorption of betamethasone 17-valerate. *Med J Austr* 1971; 1:542-544




Figure 7.—The photograph shows topical corticosteroid-induced acne rosacea.

6. Walsh P, Aeling JL, Huff L, Weston WL: Hypothalamus-pituitary-adrenal axis suppression by superpotent topical steroids. *J Am Acad Dermatol* 1993; 29:501-503
7. Feldmann RJ, Maibach HI: Regional variation in percutaneous penetration of ¹⁴C cortisol in man. *J Invest Dermatol* 1967; 48:181-183
8. Turpeinen M: Influence of age and severity of dermatitis on the percutaneous absorption of hydrocortisone in children. *Br J Dermatol* 1988; 118:517-522
9. Turpeinen M, Mashkilleysen N, Björkstén F, Salo OP: Percutaneous absorption of hydrocortisone during exacerbation and remission of atopic dermatitis in adults. *Acta Dermatol Venereol (Stockh)* 1988; 68:331-335
10. Epstein N, Epstein WL, Epstein JH: Atrophic striae in patients with inguinal intertrigo. *Arch Dermatol* 1963; 87:450
11. Stoughton RB, Cornell RL: Review of super-potent topical corticosteroids. *Semin Dermatol* 1987; 6:72-76
12. Johns AM, Bower BD: Wasting of napkin area after repeated use of fluorinated steroid ointment. *Br Med J* 1970; 1:347
13. Stenmovic DV: Corticosteroid-involved atrophy of the skin with telangiectasia: A clinical and experimental study. *Br J Dermatol* 1972; 87:548-556
14. Colomb D: Stellate spontaneous pseudoscars—Senile and presenile forms: Especially those forms caused by prolonged corticoid therapy. *Arch Dermatol* 1972; 105:551-554
15. Sibenge S, Gawkrödger DJ: Rosacea: A study of clinical patterns, blood flow, and the role of *Demodex folliculorum*. *J Am Acad Dermatol* 1992; 26:590-593
16. Plewig G, Kligman AM: Induction of acne by topical steroids. *Arch Dermatol Forsch* 1973; 247:29-52
17. Shelley WB, Shelley ED, Burmeister V: Unilateral demodectic rosacea. *J Am Acad Dermatol* 1989; 20(pt2):915-917
18. Tsuji T, Kadoya A, Tanaka R, Kono T, Hamada T: Milia induced by corticosteroids (Letter). *Arch Dermatol* 1986; 122:139-140
19. Nielsen NV, Sørensen PN: Glaucoma induced by application of corticosteroids to the periorbital region. *Arch Dermatol* 1978; 114:953-954
20. Ive FA, Marks R: Tinea incognito. *Br Med J* 1968; 3:149-152
21. Smith EB, Breneman DL, Griffith RF, et al: Double-blind comparison of naftifine cream and clotrimazole/betamethasone dipropionate cream in the treatment of tinea pedis. *J Am Acad Dermatol* 1992; 26:125-127
22. Maibach HI, Kligman AM: The biology of experimental human cutaneous moniliasis (*Candida albicans*). *Arch Dermatol* 1962; 85:113-134
23. Macmillan AL: Unusual features of scabies associated with topical fluorinated steroids. *Br J Dermatol* 1992; 87:496-497
24. Bondi EE, Kligman AM: Adverse effects from topical corticosteroids. *Proj Dermatol* 1980; 14:1-4
25. Kestel JL Jr: Hypopigmentation following the use of Cordran tape. *Arch Dermatol* 1971; 103:460
26. Wilkinson SM, Cartwright PH, English JSC: Hydrocortisone: An important cutaneous allergen. *Lancet* 1991; 337:761-762
27. Wilkinson SM, English JSC: Hydrocortisone sensitivity—Clinical features of 59 cases. *J Am Acad Dermatol* 1992; 27(pt1):683-687
28. Doooms-Goossens A: Corticosteroid contact allergy—A challenge to patch testing. *Am J Contact Dermatitis* 1993; 4:120-122
29. Doooms-Goossens A, Morren M: Results of routine patch testing with corticosteroid series in 2,073 patients. *Contact Dermatitis* 1992; 26:182-191
30. Lehtig A, Maibach HI: Immediate contact reactions, chap 6, *In* Maibach HI (Ed): *Occupational and Industrial Dermatology*, 2nd edition. Chicago, Ill, Year Book Medical Publishers, 1986, pp 32-44
31. Fisher AA, Pascher F, Vianof N: Allergic contact dermatitis due to ingredients of vehicles. *Arch Dermatol* 1968; 104:286-290
32. Vivier AD: Tachyphylaxis to topically applied steroids. *Arch Dermatol* 1976; 112:1245-1248